



# DEADFAST MOUSE & RAT KILLER PLUS SACHETS

## WHY USE THIS BAIT?

The paste sachets contain Difethialone, the most innovative active ingredient in the market presenting no known resistance and effective at a single dose.

The soft paste bait formulation has the highest palatability for enhanced performance. It outcompetes any processed, stored or prepared food.

## INSTRUCTIONS FOR USE

FOR GENERAL PUBLIC USE ONLY. USE ONLY IN TAMPER RESISTANT BAIT STATIONS.

Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it. Prior to the use of rodenticide products, non-chemical control methods (e.g. traps) should be considered.

For use indoors (mice and rats), and outdoors around buildings (rats only).

TARGET RODENT	LEVEL OF INFESTATION	INDOOR BAITING	OUTDOOR (AROUND BUILDINGS) BAITING	BEFORE BAITING	ADVICE DURING TREATMENT
<b>Brown Rat</b> <i>(Rattus norvegicus)</i>  <b>Black Rat</b> <i>(Rattus rattus)</i>	Low	Up to 150g (15 sachets) every 8 - 10 metres	Up to 150g (15 sachets) every 8 - 10 metres	Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.	The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
	High	Up to 150g (15 sachets) every 4 - 5 metres	Up to 150g (15 sachets) every 4 - 5 metres		
<b>House Mouse</b> <i>(Mus musculus)</i>	Low	Up to 50g (5 sachets) every 2 - 3 metres			
	High	Up to 50g (5 sachets) every 1 - 1.5 metres			



Bait stations should be placed in the immediate vicinity where rodent activity has been observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).

Depending on the target rodent and the level of infestation, apply the dose indicated in the table of uses and respect the distance between baiting points. Where possible, bait stations must be fixed to the ground or other structures. Place bait stations out of the reach of children, birds, pets, farm animals and other non-target animals. Place bait stations away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.

Do not place bait stations near water drainage systems where they can come into contact with water. When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product. Bait stations must be labelled with the following information: "Do not move or open"; "Contains a rodenticide"; "product name or authorisation number"; "active substance(s)", and "in case of accident call a poison centre".

Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.

Remove the remaining bait or the bait stations at the end of the treatment period.

Wear protective chemical resistant gloves during handling phase.

### FOR OUTDOOR USE

Place the bait stations in areas non-labile to flooding. Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.

### DIRECTION FOR A SAFE HANDLING AND USE

Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion. Do not use anticoagulant rodenticides as permanent baits (e.g. for prevention of rodent infestation or to detect rodent activity). Using this product should eliminate rodents within 35 days. In case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service. Search for and remove dead rodents during treatment, at least as often as baiting stations are inspected. Dispose dead rodents in accordance with local requirements.

In the UK, poisoned rodents should be double-bagged using plastic bags and either disposed of in a household waste bin with a secure lid to prevent access of wildlife or pets or collected by a specialist waste contractor or the local authority. Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

### FIRST AID INSTRUCTIONS

This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

Antidote: Vitamin K1 administered by medical/veterinary personnel only.

In case of:

- Dermal exposure, wash skin with water and then with water and soap.
- Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to an unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.

UK medical professionals should contact the National Poisons Information Service ([www.npis.org](http://www.npis.org)) for further advice.

Contact a veterinary surgeon in case of ingestion by a pet.



### DIRECTIONS FOR DISPOSAL

At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.

In the UK waste bait should be double bagged in plastic bags and disposed of in a household waste bin with a secure lid to prevent access of wildlife or pets or taken to a civic amenity site. For information on civic amenity sites contact the local authority.

### STORAGE INSTRUCTIONS

Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.

Store in places prevented from the access of children, birds, pets and farm animals.

#### DEADFAST MOUSE AND RAT KILLER PLUS SACHET

UK-2011-0003-0001 PT14 (rodenticide). For general public use only.

Contains Difethialone 0.0025 % (25 mg/kg) (CAS No 104653-34-1)



#### WARNING

May cause damage to organs (blood) through prolonged or repeated exposure. Harmful to aquatic life with long-lasting effects. Avoid release to the environment. Get medical advice if you feel unwell.

Dispose of contents/container in accordance with local regulations. Hazardous to wildlife. Dangerous for dogs and cats.

**To avoid risks to human health and the environment comply with the instructions for use**

Marketing Company: Westland Horticulture Ltd, 14 Granville Industrial Estate, Granville Road, Dungannon, Co tyrone, BT70 1NJ. Tel: +44(0) 1480 443 789